

WELLNESS PROTOCOLS FOR IMMUNE DEFENSE

Each dose is based on weight. You can split bottles if taking multiple doses. It is ok to finish the bottle if you end on a half-bottle dose. Doses should be finished within an hour. Multi-dose protocols should be taken every 12 hours approximately. If you miss a dose, resume immediately and contact us for the modified protocol as you may have to add more doses.

RECOMMENDED DOSE:

- Up to 100lbs/45kgs: 1 bottle
- 101-150lb/46-68kgs: 1.5 bottles
- 151-200lbs/69-90kgs: 2 bottles
- 201-250lbs/91-113kgs: 2.5 bottles
- 251-300+lbs/114-136kgs+: 3 bottles



CONDITION:

- **Alzheimer's** - 10 doses
- **Autism** - 2 doses (followed with 3 months of Recovery)
- **Auto-immune Diseases** (other) - Call or email for a protocol as it varies
- **C. Diff** - 8 doses (followed with Recovery for 3 months)
- **Crohn's** - 6 doses (followed with Recovery for 3 months)
- **CLL** - 4 doses
- **CMV** - 5 doses
- **Common Cold** - 1 dose (sometimes cold symptoms are actually an emotional detox and do not respond to antivirals like Immune Defense)
- **COVID -19** - 2 doses (includes those with post-Covid-19 diseases)
- **Dementia** - 2 doses
- **Diverticulitis** - 1 dose (followed by Recovery for 3 months)
- **Dysautonomia** - 3 doses
- **EBV** - 3 doses
- **Endometriosis** - 1 dose (cramping for 24-48 hours is typical after ingestion)
- **Flu Virus** - 2 doses
- **Fungal Infection** - 2 doses
- **Glaucoma** - 8 doses (followed by Immune Activator for 3+ months)
- **H. Pylori** - 2 doses (followed with Recovery for 1 month)
- **Hepatitis C** - 4 doses (warning, die-off can be significant and last several weeks)
- **Herpes breakout** - 1 dose, only if taken before lesions appear, can reduce breakout intensity
- **HIV** - 8 doses
- **HPV** - 1 dose
- **Hidradenitis Suppurativa** - 6 doses
- **Lupus** - 1 dose (it takes a few months of slow but steady improvement after ingestion)
- **Lyme and co-infections** - 16 doses
- **Malaria** - 10 doses
- **Mold Exposure** - 2 doses (and remove yourself from source)
- **Multiple Sclerosis** - 16 doses
- **Non-Hodgkin's Lymphoma** - 4 doses
- **Parkinson's** - 12 doses
- **Rheumatoid Arthritis** - 6 doses
- **Shingles** - 1 dose
- **SIBO** - 2 doses (followed with Recovery for one month)
- **Ulcerative colitis** - 11 doses (followed with Recovery for 3 months)
- **Warts** - 2 doses